


The Mean Value Theorem


Def. Let f be a real valued function defined on a metric space X . We say that f has a **local maximum** at a point $p \in X$ if there exists a $\delta > 0$ such that $f(q) \leq f(p)$ for all $q \in X$ with $d_X(p, q) < \delta$.

We say that f has a **local minimum** at a point $p \in X$ if there exists a $\delta > 0$ such that $f(p) \leq f(q)$ for all $q \in X$ with $d_X(p, q) < \delta$.



Theorem: Let $f: [a, b] \rightarrow \mathbb{R}$. If f has a local maximum or minimum at a point $p \in (a, b)$, and if $f'(p)$ exists, then $f'(p) = 0$.

Proof: Suppose $f'(p)$ exists and $f(p)$ is a local maximum.



Then by the definition of a local maximum, there exists a $\delta > 0$ such that

$$f(x) \leq f(p) \text{ for all } x \in X \text{ with } |x - p| < \delta.$$

$$|x - p| < \delta$$

$$-\delta < x - p < \delta$$

$$p - \delta < x < p + \delta.$$

Suppose that we take a point t , $p - \delta < t < p$, then we have:

$$f(t) - f(p) \leq 0 \quad \text{since } f(x) \leq f(p) \text{ for all } x \in X \text{ with } |x - p| < \delta \text{ and}$$

$$t - p < 0 \quad \text{since } t < p;$$

So we have:

$$\frac{f(t) - f(p)}{t - p} \geq 0; \text{ for } t < p. \text{ Thus we can say: } \lim_{t \rightarrow p^-} \frac{f(t) - f(p)}{t - p} \geq 0.$$

Now suppose we take a point, $p < t < p + \delta$.

So we have:

$$f(t) - f(p) \leq 0 \quad \text{since } f(x) \leq f(p) \text{ for all } x \in X \text{ with } |x - p| < \delta$$

$$t - p > 0 \quad \text{since } p < t;$$

So we have:

$$\frac{f(t) - f(p)}{t - p} \leq 0; \text{ for } t > p. \text{ That gives us: } \lim_{t \rightarrow p^+} \frac{f(t) - f(p)}{t - p} \leq 0.$$

Since $f'(p)$ exists we must have:

$$0 \leq \lim_{t \rightarrow p^-} \frac{f(t) - f(p)}{t - p} = \lim_{t \rightarrow p^+} \frac{f(t) - f(p)}{t - p} \leq 0$$

$$\text{Thus } f'(p) = \lim_{t \rightarrow p} \frac{f(t) - f(p)}{t - p} = 0.$$

A similar argument works when p is a local minimum.

The next theorem will be used later to prove L'Hopital's rule.

Theorem (Generalized Mean Value Theorem): If $f, g: [a, b] \rightarrow \mathbb{R}$, are continuous on $[a, b]$ and differentiable on (a, b) , then there exists a point $c \in (a, b)$ at which:

$$[f(b) - f(a)]g'(c) = [g(b) - g(a)]f'(c).$$

Note: we could also write this result as:

$$\frac{f(b) - f(a)}{g(b) - g(a)} = \frac{f'(c)}{g'(c)} ; \text{ This in turn could be written as:}$$

$$\frac{\frac{f(b) - f(a)}{b - a}}{\frac{g(b) - g(a)}{b - a}} = \frac{f'(c)}{g'(c)} \quad \text{i.e.}$$

$$\frac{\text{the average rate of change of } f \text{ over } [a, b]}{\text{the average rate of change of } g \text{ over } [a, b]} = \frac{\text{Inst. rate of change of } f \text{ at } c}{\text{Inst. rate of change of } g \text{ at } c}.$$

Proof: Let $h(x)$ be defined by:

$$h(x) = [f(b) - f(a)]g(x) - [g(b) - g(a)]f(x); \quad a \leq x \leq b.$$

$h(x)$ is continuous on $[a, b]$ and differentiable on (a, b) because $f(x)$ and $g(x)$ are.

Notice that $h(a) = h(b)$:

$$\begin{aligned} h(a) &= [f(b) - f(a)]g(a) - [g(b) - g(a)]f(a) \\ &= f(b)g(a) - g(b)f(a) \end{aligned}$$

$$\begin{aligned} h(b) &= [f(b) - f(a)]g(b) - [g(b) - g(a)]f(b) \\ &= -g(b)f(a) + f(b)g(a). \end{aligned}$$

If we can find a point $c \in (a, b)$ where $h'(c) = 0$ then we would have:

$$\begin{aligned} 0 &= h'(c) = [f(b) - f(a)]g'(c) - [g(b) - g(a)]f'(c) \quad \text{or} \\ [f(b) - f(a)]g'(c) &= [g(b) - g(a)]f'(c) \quad (\text{which is what we are proving}). \end{aligned}$$

So let's show we can find a point $c \in (a, b)$ where $h'(c) = 0$.

If $h(x)$ is a constant function then $h'(x) = 0$ for all $x \in (a, b)$.

If $h(x)$ is not a constant function then there is some point p , $a < p < b$ where either $h(p) > h(a)$ or $h(p) < h(a)$.

If $h(p) > h(a)$, let c be a point where h attains its global maximum (we know a continuous function on a compact set attains its absolute maximum and minimum values), $a < c < b$. This global maximum is also a local maximum because it's an interior point. Thus we know from the previous theorem that since $h(x)$ is differentiable, that $h'(c) = 0$.


Thus at this point c we have:

$$[f(b) - f(a)]g'(c) = [g(b) - g(a)]f'(c).$$

A similar argument works for $h(p) < h(a)$.

The Mean Value Theorem: Let $f: [a, b] \rightarrow \mathbb{R}$, be continuous on $[a, b]$ and differentiable on (a, b) , then there exists a c , $a < c < b$ such that:

$$\frac{f(b)-f(a)}{b-a} = f'(c).$$



Proof: Let $g(x) = x$ in the generalized mean value theorem. Then we have:

$$[f(b) - f(a)]g'(c) = [g(b) - g(a)]f'(c)$$

$$[f(b) - f(a)](1) = (b - a)f'(c)$$

$$\frac{f(b)-f(a)}{b-a} = f'(c) ; \quad a < c < b.$$

Note: The Mean Value Theorem gives us a way to bound $|f(b) - f(a)|$ for a function:

$$\frac{f(b)-f(a)}{b-a} = f'(c)$$

$$f(b) - f(a) = f'(c)(b - a)$$

$$|f(b) - f(a)| = |f'(c)|(b - a)|$$

$$\inf_{a < x < b} |f'(x)|(b - a) \leq |f(b) - f(a)| \leq \sup_{a < x < b} |f'(x)|(b - a).$$

Ex. Prove $|\sin(b) - \sin(a)| \leq |b - a|$ for all real values of a, b .

Apply the M.V.T. to $[a, b]$, for any a, b , and the function $f(x) = \sin(x)$.

$f(x) = \sin(x)$ is continuous on $[a, b]$ because it's continuous everywhere. It's differentiable on (a, b) because it's differentiable everywhere.

By the mean value theorem there exists a c , $a < c < b$ such that:

$$\frac{\sin(b) - \sin(a)}{b - a} = f'(c) = \cos(c) \quad \text{or we can write:}$$

$$\sin(b) - \sin(a) = (\cos(c))(b - a).$$

Now take absolute values:

$$|\sin(b) - \sin(a)| = |\cos(c)||b - a|; \quad \text{now use the fact that } |\cos c| \leq 1$$

$$|\sin(b) - \sin(a)| = |\cos(c)||b - a| \leq |b - a| \quad \text{so we have:}$$

$$|\sin(b) - \sin(a)| \leq |b - a| \quad \text{for all real values of } a, b.$$

Ex. Use the M. V. Theorem to prove that $\frac{1}{2} + \left(\frac{\sqrt{2}}{60}\right)\pi < \sin\left(\frac{\pi}{5}\right) < \frac{1}{2} + \left(\frac{\sqrt{3}}{60}\right)\pi$.

Apply the mean value theorem to the function $f(x) = \sin(x)$, on $\left[\frac{\pi}{6}, \frac{\pi}{5}\right]$.

Here we want to use an interval that includes $\frac{\pi}{5}$ as one endpoint and the other endpoint being a point where we "know" the value of $\sin(x)$, like $\sin\left(\frac{\pi}{6}\right)$. We could have also used $\left[\frac{\pi}{5}, \frac{\pi}{4}\right]$.

As mentioned in the previous example, $f(x) = \sin(x)$ satisfies the conditions of the mean value theorem on this interval.


By the M.V.T. we know there exists a c , $\frac{\pi}{6} < c < \frac{\pi}{5}$ such that:

$$\frac{\sin\frac{\pi}{5} - \sin\frac{\pi}{6}}{\frac{\pi}{5} - \frac{\pi}{6}} = f'(c) = \cos(c); \quad \frac{\pi}{6} < c < \frac{\pi}{5}$$

$$\frac{\sin\frac{\pi}{5} - \frac{1}{2}}{\frac{\pi}{30}} = \cos(c) \quad \frac{\pi}{6} < c < \frac{\pi}{5};$$

since $\cos(x)$ is decreasing on $[\frac{\pi}{6}, \frac{\pi}{4}]$

$$\frac{\sqrt{3}}{2} = \cos\frac{\pi}{6} > \cos(c) > \cos\frac{\pi}{5} > \cos\frac{\pi}{4} = \frac{\sqrt{2}}{2};$$



$$\frac{\sqrt{3}}{2} > \frac{\sin\frac{\pi}{5} - \frac{1}{2}}{\frac{\pi}{30}} > \frac{\sqrt{2}}{2} \quad \text{now let's solve the inequality for } \sin\frac{\pi}{5}$$

$$\frac{\sqrt{3}}{2} \left(\frac{\pi}{30}\right) > \sin\left(\frac{\pi}{5}\right) - \frac{1}{2} > \frac{\sqrt{2}}{2} \left(\frac{\pi}{30}\right)$$

$$\frac{1}{2} + \frac{\sqrt{3}}{2} \left(\frac{\pi}{30}\right) > \sin\left(\frac{\pi}{5}\right) > \frac{1}{2} + \frac{\sqrt{2}}{2} \left(\frac{\pi}{30}\right) \quad \text{or}$$

$$\frac{1}{2} + \left(\frac{\sqrt{2}}{60}\right)\pi < \sin\left(\frac{\pi}{5}\right) < \frac{1}{2} + \left(\frac{\sqrt{3}}{60}\right)\pi.$$

Ex. Let $f(x) = \tan^{-1}x$ and apply the M.V.T. to $[a, b]$; $a, b > 0$ to prove:

a.
$$\frac{b-a}{1+b^2} < \tan^{-1}b - \tan^{-1}a < \frac{b-a}{1+a^2} .$$

b. Apply part "a" to $[1, \frac{4}{3}]$ to get the inequality:

$$\frac{\pi}{4} + \frac{3}{25} < \tan^{-1}\left(\frac{4}{3}\right) < \frac{\pi}{4} + \frac{1}{6} .$$

a. $f(x) = \tan^{-1}x$ is continuous everywhere and differentiable everywhere and thus it satisfies the M.V. T. on any interval $[a, b]$.

By the M.V.T. there exists a c , $0 < a < c < b$ such that:

$$\frac{\tan^{-1}b - \tan^{-1}a}{b-a} = \frac{1}{1+c^2}; \quad 0 < a < c < b \quad (\text{since } f'(x) = \frac{1}{1+x^2}).$$

Since $0 < a < c < b$ we know that $a^2 < c^2 < b^2$ and $1 + a^2 < 1 + c^2 < 1 + b^2$,

and finally $\frac{1}{1+a^2} > \frac{1}{1+c^2} > \frac{1}{1+b^2}$.

Now since $\frac{\tan^{-1}b - \tan^{-1}a}{b-a} = \frac{1}{1+c^2}$, replacing in the above inequality we get:

$$\frac{1}{1+a^2} > \frac{\tan^{-1}b - \tan^{-1}a}{b-a} > \frac{1}{1+b^2} .$$

Now multiply through by $(b - a)$, which is positive because $b > a$:

$$\frac{b-a}{1+b^2} < \tan^{-1}b - \tan^{-1}a < \frac{b-a}{1+a^2}.$$

b. Applying this inequality when $a = 1$ and $b = \frac{4}{3}$

$$\frac{\frac{4}{3}-1}{1+(\frac{4}{3})^2} < \tan^{-1}\left(\frac{4}{3}\right) - \tan^{-1}1 < \frac{\frac{4}{3}-1}{1+(1)^2}$$

$$\frac{\frac{1}{3}}{\frac{25}{9}} < \tan^{-1}\left(\frac{4}{3}\right) - \frac{\pi}{4} < \frac{1}{6}$$

$$\frac{\pi}{4} + \frac{3}{25} < \tan^{-1}\left(\frac{4}{3}\right) < \frac{\pi}{4} + \frac{1}{6}.$$

Ex. Prove that $e^x > 1 + x$ for $x > 0$.

Apply the M.V.T. to the function $f(x) = e^x$ on the interval $[0, x]$.

$f(x)$ satisfies the M.V.T. because it's continuous everywhere and differentiable everywhere.

By the M.V.T. we know that there is a c , $0 < c < x$ such that

$$\frac{e^x - e^0}{x - 0} = f'(c) = e^c; \quad 0 < c < x \quad \text{or}$$

$$\frac{e^x - 1}{x} = e^c; \quad 0 < c < x.$$

Since $0 < c$ and $f(x) = e^x$ is an increasing function $e^0 = 1 < e^c$. Thus we have:

$$\frac{e^x - 1}{x} = e^c > 1; \quad \text{Now solve this inequality for } e^x.$$

$$e^x - 1 > x$$

$$e^x > 1 + x \quad \text{for } x > 0.$$

Ex. Suppose $f'(x)$ exists on (a, b) and $\sup_{a < x < b} |f'(x)| \leq M$, show that $f(x)$ is uniformly continuous on (a, b) .

Let $x, y \in (a, b)$ then by the M.V.T. we have:

$$|f(x) - f(y)| \leq M|x - y|.$$

So if we choose $\delta = \epsilon/M$ then:

$$|f(x) - f(y)| \leq M|x - y| < M(\delta) = M\left(\frac{\epsilon}{M}\right) = \epsilon$$

and f is uniformly continuous on (a, b) .

Def. A function $f: \mathbb{R} \rightarrow \mathbb{R}$ is called **monotonically increasing** if $p > q$ implies that $f(p) \geq f(q)$. f is called **monotonically decreasing** if $p > q$ implies that

$$f(p) \leq f(q).$$



Theorem: Suppose f is differentiable in (a, b) :

- If $f'(x) \geq 0$ for all $x \in (a, b)$ then f is monotonically increasing.
- If $f'(x) = 0$ for all $x \in (a, b)$ then f is a constant function
- If $f'(x) \leq 0$ for all $x \in (a, b)$ then f is monotonically decreasing.

Proof: Take any two point $p, q \in (a, b)$ with $p > q$. Since f is differentiable in (a, b) , it is also differentiable in (q, p) and continuous in $[q, p]$ (if f is differentiable at a point t , then it is continuous at t) thus f satisfies the conditions of the Mean Value Theorem on $[q, p]$.

Thus we know:

$$f(p) - f(q) = f'(c)(p - q) \quad \text{where } q < c < p.$$

a. If $f'(x) \geq 0$ for all $x \in (a, b)$ then $f'(c) \geq 0$, and thus $f'(c)(p - q) \geq 0$;

Hence $f(p) - f(q) = f'(c)(p - q) \geq 0$ and $f(p) \geq f(q)$.

So f is monotonically increasing.

b. If $f'(x) = 0$ for all $x \in (a, b)$ then $f'(c) = 0$, and thus $f(p) - f(q) = 0$;

or $f(p) = f(q)$. So f is a constant function.

c. If $f'(x) \leq 0$ for all $x \in (a, b)$ then $f'(c) \leq 0$, and thus $f'(c)(p - q) \leq 0$;

Hence $f(p) - f(q) = f'(c)(p - q) \leq 0$ and $f(p) \leq f(q)$.

So f is monotonically decreasing.

Ex. Suppose f is differentiable everywhere and $f(2) = 6$ and $|f'(x)| \leq 4$, for all values of x . show that $-6 \leq f(5) \leq 18$ and $-2 \leq f(0) \leq 14$.

Since f is differentiable everywhere it satisfies the Mean Value Theorem on any closed interval $[a, b]$. If we apply the M.V.T. to the interval $[2, 5]$ we get:

$$f(5) - f(2) = f'(c)(5 - 2) \quad \text{where } 2 < c < 5;$$

Since $f(2) = 6$, we have:

$$f(5) - 6 = (f'(c))(3).$$

Since $|f'(x)| \leq 4$, we know that $-4 \leq f'(x) \leq 4$ for all x , so
 $-4 \leq f'(c) \leq 4$

and $-12 \leq (f'(c))(3) \leq 12$.

Since $f(5) - 6 = (f'(c))(3)$ we have:

$$-12 \leq f(5) - 6 \leq 12 \quad \text{or}$$

$$-6 \leq f(5) \leq 18 .$$

Now let's apply the M.V.T. to the interval $[0,2]$

$$f(2) - f(0) = f'(c)(2 - 0) \quad \text{where } 0 < c < 2;$$

Since $f(2) = 6$, we have:

$$6 - f(0) = (f'(c))(2).$$

Since $|f'(x)| \leq 4$, we know that $-4 \leq f'(x) \leq 4$ for all x , so

$$-4 \leq f'(c) \leq 4$$

$$\text{and } -8 \leq (f'(c))(2) \leq 8 .$$

Since $6 - f(0) = (f'(c))(2)$ we have:

$$-8 \leq 6 - f(0) \leq 8$$

$$-14 \leq -f(0) \leq 2$$

$$14 \geq f(0) \geq -2 .$$

In fact, if $f(x)$ satisfies the Mean Value Theorem on an interval, and $L \leq f'(x) \leq K$ on that interval then we have:

$$L \leq \frac{f(x)-f(a)}{x-a} = f'(c) \leq K.$$

Solving this inequality for $f(x)$ we get:


$$f(a) + L(x - a) \leq f(x) \leq f(a) + K(x - a) \quad \text{if } x \geq a$$

$$f(a) + L(x - a) \geq f(x) \geq f(a) + K(x - a) \quad \text{if } x \leq a.$$

Which means the values of $f(x)$ can't go outside the lines

$$y = f(a) + L(x - a) \quad \text{and} \quad y = f(a) + K(x - a)$$

on the interval.



Ex. Suppose f is differentiable everywhere and $f(1) = 7$ and $f'(x) \geq -3$, for all values of x . Show that $f(6) \geq -8$. Can we find an upper bound on $f(6)$?

Since f is differentiable everywhere it satisfies the Mean Value Theorem on any closed interval $[a, b]$. If we apply the M.V.T. to the interval $[1, 6]$ we get:

$$f(6) - f(1) = f'(c)(6 - 1) \quad \text{where } 1 < c < 6.$$

Since $f(1) = 7$, we have:

$$f(6) - 7 = f'(c)(5) \quad \text{where } 1 < c < 6.$$

Since $f'(x) \geq -3$ for all values of x , we know that $f'(c) \geq -3$.

Thus we have:

$$f(6) - 7 = f'(c)(5) \geq (-3)(5) = -15; \quad \text{now add 7 to both sides}$$
$$f(6) \geq -8.$$

We cannot find an upper bound on $f(6)$ because we have no upper bound on $f'(c)$.

Theorem (L'Hopital's rule) Suppose f, g are real valued differentiable functions on (a, b) , and $g'(x) \neq 0$ for all $x \in (a, b)$, where $-\infty \leq a, b \leq +\infty$. Suppose $p \in (a, b)$ (so p could be either $+\infty$ or $-\infty$) and $\lim_{x \rightarrow p} \frac{f'(x)}{g'(x)} = A$.

If $\lim_{x \rightarrow p} f(x) = 0$, $\lim_{x \rightarrow p} g(x) = 0$, or

If $\lim_{x \rightarrow p} f(x) = \pm\infty$, $\lim_{x \rightarrow p} g(x) = \pm\infty$

Then $\lim_{x \rightarrow p} \frac{f(x)}{g(x)} = \lim_{x \rightarrow p} \frac{f'(x)}{g'(x)} = A$.

Proof: We'll just prove the case where $\lim_{x \rightarrow p} f(x) = \lim_{x \rightarrow p} g(x) = 0$ and $p \neq \pm\infty$.

Since $\lim_{x \rightarrow p} f(x) = \lim_{x \rightarrow p} g(x) = 0$ and both f, g are continuous at $x = p$,

$$f(p) = g(p) = 0.$$

Choose an $x > p$. Since f, g are differentiable everywhere, they satisfy the Extended Mean Value Theorem on $[p, x]$, so we can conclude that:

$$\frac{f(x) - f(p)}{g(x) - g(p)} = \frac{f'(c)}{g'(c)} \quad \text{for } p < c < x.$$

Since $f(p) = g(p) = 0$ we have:

$$\frac{f(x)}{g(x)} = \frac{f'(c)}{g'(c)} \quad \text{for } p < c < x.$$

Now take the limit from the right:

$$\lim_{x \rightarrow p^+} \frac{f(x)}{g(x)} = \lim_{x \rightarrow p^+} \frac{f'(c)}{g'(c)} = \lim_{x \rightarrow p^+} \frac{f'(x)}{g'(x)}.$$

Now Choose an $x < p$. Since f, g are differentiable everywhere, they satisfy the Extended Mean Value Theorem on $[x, p]$, so we can conclude that:

$$\frac{f(p) - f(x)}{g(p) - g(x)} = \frac{f'(c)}{g'(c)} \quad \text{for } x < c < p.$$

Since $f(p) = g(p) = 0$ we have:

$$\frac{f(x)}{g(x)} = \frac{f'(c)}{g'(c)} \quad \text{for } x < c < p.$$

Now take the limit from the left:

$$\lim_{x \rightarrow p^-} \frac{f(x)}{g(x)} = \lim_{x \rightarrow p^-} \frac{f'(c)}{g'(c)} = \lim_{x \rightarrow p^-} \frac{f'(x)}{g'(x)}.$$

Since, by assumption $\lim_{x \rightarrow p} \frac{f'(x)}{g'(x)} = A$, the right hand and left hand limits must be the same (for both $\frac{f'(x)}{g'(x)}$ and $\frac{f(x)}{g(x)}$).

Thus we have: $\lim_{x \rightarrow p} \frac{f(x)}{g(x)} = \lim_{x \rightarrow p} \frac{f'(x)}{g'(x)} = A$.

Ex. Find $\lim_{x \rightarrow 2} \frac{x^2 - 2x}{\sin(x-2)}$.

$\lim_{x \rightarrow 2} (x^2 - 2x) = 0$, and $\lim_{x \rightarrow 2} \sin(x - 2) = 0$, so by L'Hopital's rule:

$$\lim_{x \rightarrow 2} \frac{x^2 - 2x}{\sin(x-2)} = \lim_{x \rightarrow 2} \frac{2x-2}{\cos(x-2)} = \frac{2}{1} = 2.$$