

## Changing Bases- HW Problems

For all of the following problems when you are changing bases use a change of basis matrix  $P$ .

1. Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  by  $T(x_1, x_2) = (2x_1 + x_2, x_1 - x_2)$ .
  - a. Find a matrix  $A$  which represent  $T$  in the standard basis  $\{\vec{e}_1, \vec{e}_2\}$  for  $\mathbb{R}^2$ .
  - b. Let  $v_1 = e_1 + e_2$  and  $v_2 = e_1 - e_2$ .  $\{v_1, v_2\}$  is another basis for  $\mathbb{R}^2$ . Find a matrix representation,  $B$ , of  $T$  with respect to  $\{v_1, v_2\}$  (for both  $\mathbb{R}^2$ 's).
  - c. Let  $w_1 = -e_1 + 3e_2$  and  $w_2 = 2e_1 - e_2$ . Find a matrix representation,  $C$ , of  $T$  with respect to  $\{w_1, w_2\}$  (for both  $\mathbb{R}^2$ 's).
  - d. Show that using the matrices from parts b and c you can find matrix  $C$  from matrix  $B$  and the change of basis matrix from  $\{w_1, w_2\}$  to  $\{v_1, v_2\}$ .

In problems 2-5  $A$  is a matrix representation of a linear transformation in the standard basis. Find a matrix representation of the linear transformation in the new basis.

$$2. \quad A = \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}; \quad \text{new basis} = \{ \langle 1, 2 \rangle, \langle 1, 1 \rangle \}$$

$$3. \quad A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}; \quad \text{new basis} = \{ \langle 7, 3 \rangle, \langle 2, 1 \rangle \}$$

$$4. \quad A = \begin{bmatrix} 1 & 0 & 2 \\ -1 & 1 & 3 \\ 0 & 1 & 1 \end{bmatrix};$$

$$\text{new basis} = \{ \langle 1, 1, 1 \rangle, \langle 2, 1, 0 \rangle, \langle -1, 1, 1 \rangle \}$$

$$\text{You can assume that if } P = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix} \text{ then } P^{-1} = \frac{1}{2} \begin{bmatrix} 1 & -2 & 3 \\ 0 & 2 & -2 \\ -1 & 2 & -1 \end{bmatrix}$$

$$5. \quad A = \begin{bmatrix} 2 & 1 & -1 \\ 0 & 1 & 3 \\ -1 & 2 & 1 \end{bmatrix};$$

$$\text{new basis} = \{ \langle 0, -2, 1 \rangle, \langle 1, 2, 0 \rangle, \langle 1, 1, 1 \rangle \}.$$

$$\text{You can assume that if } P = \begin{bmatrix} 0 & 1 & 1 \\ -2 & 2 & 1 \\ 1 & 0 & 1 \end{bmatrix} \text{ then } P^{-1} = \begin{bmatrix} 2 & -1 & -1 \\ 3 & -1 & -2 \\ -2 & 1 & 2 \end{bmatrix}$$

6. Suppose  $T: P_2(\mathbb{R}) \rightarrow P_2(\mathbb{R})$  by

$$T(a_0 + a_1x + a_2x^2) = (a_0 + a_1) + (a_2 - a_1)x + (a_2 - a_0)x^2.$$

a. Find a matrix representation of  $T$  with respect to the standard basis for  $P_2(\mathbb{R})$ ,  $\{1, x, x^2\}$ .

b. Find a matrix representation of  $T$  with respect to the basis  $\{1 + x, x + x^2, 1 + x^2\}$ .

You can assume that if  $P = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$  then  $P^{-1} = \frac{1}{2} \begin{bmatrix} 1 & 1 & -1 \\ -1 & 1 & 1 \\ 1 & -1 & 1 \end{bmatrix}$ .

c. Find a matrix representation of  $T$  with respect to the basis  $\{1, 1 + x, 1 + x + x^2\}$ .

You can assume that if  $P = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$  then  $P^{-1} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}$ .

7.  $n \times n$  matrices  $A$  and  $B$  are similar if there exists an invertible  $n \times n$  matrix  $Q$  such that  $B = Q^{-1}AQ$ . Show that if  $A$  and  $B$  are similar matrices then  $\det(A) = \det(B)$ .

8. Suppose that  $A, B$ , and  $C$  are  $n \times n$  matrices. Show

a.  $A$  is similar to  $A$

b. If  $A$  is similar to  $B$ , then  $B$  is similar to  $A$ .

c. If  $A$  is similar to  $B$ , and  $B$  is similar to  $C$ , then  $A$  is similar to  $C$ .

Note: a, b, and c mean that similarity of matrices is an equivalence relation.

9. Suppose that  $A$  is an invertible  $n \times n$  matrix and  $A$  is similar to  $B$ .
- Prove that  $B$  is invertible.
  - Prove that  $A^{-1}$  is similar to  $B^{-1}$ .